EPD™

Environmental Product Declaration

Global GreenTag EPD Program:

Compliant to EN15804+A2 2019

Xypex Chemical Corporation

Xypex Patch'n Plug

13731 Mayfield Place
Richmond British Columbia
Canada

XYPEX®
Concrete Solutions

Mandatory Disclosures

EPD type	Cradle to grave A1 to C4 + D	**EPD Numbers**	XYP042022EP
Issue Date	07 March 2023	**Valid Until**	07 March 2028

Demonstration of Verification

PCR	Standard EN 15804+A2 2019 serves as core Product Category Rules (PCR) [1]. Sub-PCR UCM:2023 Unreinforced Concrete Mixtures and Additives also applies [2].
☑ Internal	Delwyn Jones 02 Feb 2023 — LCA Developed by Delwyn Jones, The Evah Institute 08Feb2023 — LCA Reviewed by Direshni Naiker The Evah Institute 07/03/2023 — EPD Reviewed by David Baggs, Global GreenTag Pty Ltd
☑ External	08Feb2023 — Third Party Verifier[a] Mathilde Vlieg Malaika LCT a. Independent external verification of the declaration and data, mandatory for business-to-consumer communication according to ISO 14025:2010 [2].
Communication	This EPD discloses potential environmental outcomes compliant with EN 15804 for business-to-business communication.
Comparability	Construction product EPDs may not be comparable if not EN15804 compliant. Different program EPDs may not be comparable. Comparability is further dependent on the product category rules and data source used.
Reliability	LCIA results are relative expressions that do not predict impacts on category endpoints, exceeding of thresholds, safety margins or risks.
Owner	This EPD is the property of the declared manufacturer.
Explanations	Further explanatory information is available at info@globalgreentag.com or by contacting certification1@globalgreentag.com [3].

EPD Program Operator	LCA and EPD Producer	Declaration Owner
Global GreenTag Pty Ltd	Ecquate Pty Ltd	Xypex Chemical Corporation
PO Box 311 Cannon Hill	PO Box 123 Thirroul	13731 Mayfield Place,
QLD 4170 Australia	NSW 2515 Australia	Richmond BC Canada
Phone: +61 (0)7 33 999 686	Phone: +61 (0)7 5545 0998	Phone: +1 604.273.5265
http://www.globalgreentag.com	http://www.evah.com.au	https://www.xypex.com/

Program Description

EPD type	Cradle to grave A1 to C4 + D as defined by EN 15804 [1]
System boundary	The system boundary with nature includes material and energy acquisition, processing, manufacture, transport, installation, use plus waste arising.to end of life.
Stages included	Stages A1-3 A4-5, B1-4, C1 to C2 and C4 D1 to D3
Stages excluded	No stage was excluded but B1, B2, B4 to B7 and C3 flows were all zero.
Scope Depiction	Figure 1 depicts all modules being declared including some with zero results. Any module not declared (MND) does not indicate a zero result.

Model	Actual					Scenarios											Potential		
Information	Building Life Cycle Assessment																Supplementary		
Stages	Product			Construct		Use							End-of-Life				Benefit & load beyond system		
Data Modules						Fabric					Operation								
Unit Operations	A1	A2	A3	A4	A5	B1	B2	B3	B4	B5	B6	B7	C1	C2	C3	C4	D1	D2	D3
Cradle to Gate+ Options & Grave	Resource	Transport	Manufact-ure	Transport	Construct	Use	Maintain	Repair	Replace	Refurbish	Energy use	Water use	Demolish	Transport	Process Waste	Disposal	Reuse	Recovery	Recycling

Figure 1 EPD Life Cycle Modules Cradle to Grave

Data Sources

Primary Data	Data was collected from primary sources 2019 to 2022 including the manufacturer and suppliers' standards, locations, logistics, technology, market share, management system in accordance with EN ISO 14044:2006, 4.3.2, [4]. All are biochemical-physical allocated none are economically allocated.
A1-A3 Stage inclusions	Operations include all known raw material acquisition, refining and processing plus scrap or material reuse from prior systems; electricity generated from all sources with extraction, refining & transport plus secondary fuel energy and recovery processes. Also, transport to factory gate; manufacture of inputs, ancillary material, product, packaging, maintenance, replacement plus flows leaving at end-of-waste boundary and fates of all flows at end of life.
Variability	Significant differences of average LCIA results are declared.
Chemicals of Concern	Contains no substances in the European Chemicals Agency "Authorised or Candidate Lists of Substances of Very High Concern (SVHCs)".

Data Quality

Data cut-off & quality criteria complies with EN 15804 [1] The LCA used background data aged <10 years and quality parameters tabled below.

Background	**Data Quality**	**Parameters and Uncertainty (U)**			
Correlation	**Metric σg**	U ±0.01	U ±0.05	U ±0.10	U ±0.20
Reliability	**Reporting**	Site Audit	Expert verify	Region	Sector
	Sample	>66% trend	>25% trend	>10% batch	>5% batch
Completion	**Including**	>50%	>25%	>10%	>5%
	Cut-off	0.01%w/w	0.05%w/w	0.1%w/w	0.5%w/w
Temporal	**Data Age**	<3 years	≤5 years	<7.5 years	<10 years
	Duration	>3 years	<3 years	<2 years	1 year
Technology	**Typology**	Actual	Comparable	In Class	Convention
Geography	**Focus**	Process	Line	Plant	Corporate
	Range	Continent	Nation	Plant	Line
	Jurisdiction	Representation is Global. Africa, North America, Europe, Pacific Rim			

Product Information

A fast-setting, hydraulic cement compound for concrete patching and repair to seal cracks, tie holes, and other defects. It stops flowing water in seconds with sealing enhanced by its crystalline waterproofing.

Brand Name & Code	Patch'n'Plug	**Range Names**	Xypex Patch'n'Plug
Factory warranty	One year	**Reference Service Life**	60 years [5,6]
Manufacturer	Xypex Chemical Corporation		
Manufacturer address	13731 Mayfield Place, Richmond British Columbia, Canada		
Site representation	Canadian and American		
Function in Building	Repair mortar for patching and resurfacing of deteriorated concrete		
Functional unit	Cradle to grave concrete repair, remedial & waterproofing/kg 60years		

Manufacturer Safety Data Sheets inform about user protection as the alkalinity may irritate skin and eyes.

Safety Procedures	https://www.xypex.com/technical/safety-data
Specifications	https://www.xypex.com/technical/specifications
Practices References	https://www.xypex.com/technical/statements
Installation Procedures	https://www.xypex.com/products/installations

Product Components

This section summarises factory components, functions, source nation and mass share. In product content listed below the % dry mass has a ±5% range and a confidence interval that is 90% certain to contain true population means at any time. Listing such 90±5% certainty considers normal resource acquisition, supply chain, sedimentation, seasonal, manufacturing and product variation over this EPD's validity period. This also allows for intellectual property protection and fullest possible transparency.

Function	Component	Cradle	Amount
Cement Binder	Cement	Canada	>30 <40
Aggregate	Moraine sand	Canada	>25 <35
Cement Binder	Calcium Aluminate	Virginia	>15 <25
Crystalline Waterproofing	Base mix	Canada	>5 <15
Hydration	Hydrated Lime	Canada	>5 <15
Packaging			
Pallet wood	Wood	Canada	>1.5 <2.0
Pail, Straps, Wrap & Tape	Polymers	Canada	>0.4 <0.5
Packaging	Cardboard and paper	Canada	>0.2 <0.3

Product Functional & Technical Performance Information

This section provides specifications and data to calculate results factoring different mass and period. Small and large pail dry power and wet mortar volume capacities are listed below.

Pail Capacity	Small	Mortar	Large	Mortar
Measure	**dry mass**	**wet volume**	**dry mass**	**wet volume**
Imperial System	20 lb	0.18 ft^3	60 lb	0.54 ft^3
Metric System	9.07 kg	0.0051 m^3	27.2 kg	0.0154 m^3

Laboratory test samples use 1 to 3.25 volumes water to dry powder. Results may vary with field conditions.

Performance	Test Method	Conformance Results		Period
Setting Time	ASTM C 266	1:30 to 4.0 minutes:seconds		Initial
		4:30 to 9:0 minutes:seconds		Final
Compressive Strength	ASTM C 109	1740 psi	1 day	12 MPa
		3630 psi	28 days	25 MPa
Tensile Bond Pull-off	CSA A23.2-6B	120psi		0.8MPa

System Analysis Scope and Boundaries

Stages A1 to 3 model actual operations. Stage A4 to C4 are model scenarios.

Typical scenarios are assumed to forecast unit operations as described in the next section.

Figure 2. shows included processes in a cradle to grave system boundary to end of life fates to unshown beyond the boundary reuse, recycling or landfill grave.



Figure 2. Product Process Flow Chart Completeness

Scenarios for Modules (Units/Functional Unit)

This section defines modelling scenarios beyond actual A1 to A3 operations from stage A4 to D3.

A Construction	Type specified	Amount	Type specified	Amount
A4 Transport to Site	25t semi-trailer	60 km	85% Capacity	Full back load
Volume capacity (<1 to ≥1)	Utilisation factor	1	Uncompressed	Un-nested
A5 Installation utilities	Town water	0.53 litre	Grid power	0.0002 MJ
Waste on site	Spill	0.05kg		
Scrap collection & routes	25t semi-trailer	60 km	to landfill	In LCA report

Stage B2 and B3 scenarios are listed below. Stages B1 Use of building fabric, B4 Replacement, B5 Refurbishment, B6 Building Operating Energy and B7 Building Operating Water all have zero flows.

B Building	Type specified	Amount	Type specified	Amount
B2 Maintenance	None typical	nil	Clean cycle	nil
B3 Repair 5%	As per website	Specified	Freight to site	As A5

Stage C1, C2 and C4 scenarios are listed below. Stage C3 Waste Treatment has zero flows.

C End of Life	Type specified	Amount	Type specified	Amount
C1 Demolition	Remove worn area	0.01kg	Collect separately	0.01g
C2 Transport	25t truck road	50km	85% capacity	No back load
C4 Disposal	Product specific	0.01kg	Collect separately	0.01kg
Recovery system	No recycling	0.0 kg	Not for energy	0.0 kg

Stage D scenarios D1 Reuse and D2 Recovery are listed below. D3 Recycling has zero flows.

D Beyond System Boundary	Type specified	Amount	Type specified	Amount
D1 Reuse	typically all	100%	typically all	1.00kg
D2 Recovery	typically all	100%	Cleaning	sweep
D3 Recycle	At 60 years	Nil	Typically none	0%

Environmental Impact Terminology

Environmental impacts contributing to risks of social and ecological issues and collapse are tabled below with common names and remedies given for each indicator.

Global warming forcing Climate Change	Greenhouse gases absorb infra-red radiation. This heat reduces thermal energy differentials, from equator to poles, forcing ocean current and wind circulation to blend and regulate climate. Weakly blended "lumpier" weather has more frequent, extreme heat wave, fire-storm, cyclone, rain-storm, flood and blizzard events. Accumulation of carbon dioxide, natural gas methane, nitrous oxides and volatile organic compounds from burning fossil fuels causes global warming. Forest and wilderness growth absorbing air-borne carbon in biomass can drawdown such accumulation. Urgent renewable energy reliance is vital in time to avoid imminent tipping points and the worsening "***climate emergency***".
Ozone layer depletion	Stratospheric ozone loss weakens the planet's solar shield so more shorter wavelength ultraviolet (UVB) light reaching earth damages plants and increases malignant melanoma and skin cancer in humans and animals. Chlorofluorocarbons, hydrochlorofluorocarbons (HCFC), chlorobromomethane, hydrobromofluorocarbons, carbon tetrachloride, methyl chloroform, methyl bromide and halon gas cause ozone layer loss. To repair the "***ozone hole***" reliance on ozone-safe refrigerants, aerosols and solvents is essential to avoid further its depletion and enable accumulation of naturally-formed ozone.
Acidification	Acidification reduces soil and waterway pH, impedes nitrogen fixation vital for plant growth and inhibits natural decomposition. It increases rates and incidence of fish kills, forest loss and deterioration of buildings and materials. Chief synthetic causes of "***acid rain***" are emissions of sulphur and nitrogen oxides, hydrochloric and hydrofluoric acids and ammonia from burning fossil fuels polluting precipitation of rain and snow world-wide.
Eutrophication of terrestrial, freshwater and marine life	Eutrophication from excessively high macronutrient levels added to natural waters promotes excessive plant growth that severely reduces oxygen, water and habitat security for aquatic and terrestrial organisms across related ecosystems. Chief synthetic cause of "***algal blooms***" is nitrogen (N, NOx, NH_4) and phosphorus (P, PO_4^{3-}) in rain run-off over-fertilised land catchments.
Photochemical ozone creation	Tropospheric photochemical ozone, called "***summer smog***" near ground level, is created from natural and synthetic compounds in UV sunlight. Low concentration smog damages vegetation and crops. High concentration smog is hazardous to human health. Chief synthetic causes are nitrogen oxides, carbon monoxide and volatile organic compounds (VOC) pollutants. Avoiding reliance on dirtiest coal fuel and volatile chemicals has reduced smog incidence in many areas globally.
Depletion of minerals, metals & water	Abiotic depletion of finite mineral resources increases time, effort and money required to obtain more resources to the point of extinction of naturally viable reserves. This can limit access to available, valuable and scarce elements vital for human-life. The youth movement "***extinction rebellion***" calls on adults to secure climate, reserves and biodiversity for current and future generations.
Depletion of fossil fuel reserves	Abiotic depletion of resources by consuming finite oil, natural gas, coal and yellowcake fossil fuel reserves leaves current and future generations suffering limited available, accessible, plentiful, essential valuable as well as scarce raw material, medicinal, chemical, feedstock and fuel stock. Approaching "***peak oil***" acknowledged fossil fuel reserves are finite and the need for decision-makers to act to avoid market instability, insecurity and or oil and gas wars.

Glossary of Terms, Methods and Units

Acronyms, methods and units of impact potentials plus inventory inputs and outputs, are defined below

Impact Potentials	Acronym	Description of Methods	Units
Climate Change fossil	GWP_{ff}	GWP fossil fuels [7]	kg CO_{2eq}
Climate Change biogenic	GWP_{bio}	GWP biogenic [7]	kg CO_{2eq}
Climate Change land use	GWP_{luluc}	GWP land use & change [7]	kg CO_{2eq}
Climate Change total	GWP_{t}	Global Warming Potential [7]	kg CO_{2eq}
Stratospheric Ozone Depletion	ODP	Stratospheric Ozone Loss [8]	kg CFC_{11eq}
Photochemical Ozone Creation	POCP	Summer Smog [9]	kg $NMOC_{eq}$
Acidification Potential	AP	Accumulated Exceedance [10]	mol H^{+}_{eq}
Eutrophication Freshwater	EP_{fresh}	Excess nutrients freshwater [11]	kg P_{eq}
Eutrophication Marine	EP_{marine}	Excess marine nutrients [11]	kg N_{eq}
Eutrophication Terrestrial	EP_{land}	Excess Terrestrial nutrients [11]	mol N_{eq}
Mineral & Metal Depletion	ADP_{min}	Abiotic Depletion minerals [12]	kg Sb_{eq}
Fossil Fuel Depletion	ADP_{ff}	Abiotic Depletion fossil fuel [13]	MJ $_{ncv}$
Water Depletion	WDP	Water Deprivation Scarcity [14, 15]	m^3 $_{WDP\ eq}$
Fresh Water Net	FW	Lake, river, well & town water	m^3
Secondary Material	SM	Post-consumer recycled (PCR)	kg
Secondary Renewable Fuel	RSF	PCR biomass burnt	MJ $_{ncv}$
Primary Energy Renewable Material	PERM	Biomass retained material	MJ $_{ncv}$
Primary Energy Renewable Not Feedstock	PERE	biomass fuels burnt	MJ $_{ncv}$
Primary Energy Renewable Total	PERT	Biomass burnt + retained	MJ $_{ncv}$
Secondary Non-renewable Fuel	NRSF	PCR fossil-fuels burnt	MJ $_{ncv}$
Primary Energy Non-renewable Material	PENRM	Fossil feedstock retained	MJ $_{ncv}$
Primary Energy Non-renewable Not Feedstock	PENRE	fossil-fuel used or burnt	MJ $_{ncv}$
Primary Energy Non-renewable Total	PENRT	Fossil feedstock & fuel use	MJ $_{ncv}$
Hazardous Waste Disposed	HWD	Reprocessed to contain risks	kg
Non-hazardous Waste Disposed	NHWD	Municipal landfill facility waste	kg
Radioactive Waste Disposed	RWD	Mostly ex nuclear power stations	kg
Components For Reuse	CRU	Product scrap for reuse as is	kg
Material For Recycling	MFR	Factory scrap to remanufacture	kg
Material For Energy Recovery	MER	Factory scrap use as fuel	kg
Exported Energy Electrical	EEE	Uncommon for building products	MJ $_{ncv}$
Exported Energy Thermal	EET	Uncommon for building products	MJ $_{ncv}$

Results Module A: Cradle to Site

Table 1 shows results for A1 -A3 Acquisition, Transport and Manufacture then A4 Delivery and A5 Construct

Table 1 A1 to A5 Impact & Inventory Results/Functional Unit

Result	A1-3	A4	A5
Climate Change biogenic	-5.1E-03	-1.0E-06	-4.3E-04
Climate Change luluc (landuse)	3.9E-06	1.7E-09	1.7E-07
Climate Change fossil	1.3	1.9E-02	5.1E-02
Climate Change total	1.3	1.9E-02	5.1E-02
Stratospheric Ozone Depletion	1.9E-08	1.7E-13	8.0E-10
Photochemical Ozone Creation	5.0E-03	1.2E-04	2.3E-04
Acidification Potential	2.3E-03	1.2E-05	1.0E-04
Eutrophication Freshwater	5.9E-08	5.6E-10	9.8E-09
Eutrophication Marine	5.4E-04	2.3E-06	2.4E-05
Eutrophication Terrestrial	1.3E-03	7.9E-06	5.8E-05
Fossil Depletion	0.57	2.3E-02	2.4E-02
Mineral and Metal Depletion	4.6E-04	7.2E-06	1.7E-05
Water Scarcity Depletion	1.4E-02	3.0E-06	5.8E-04
Net Fresh Water Use	88	0.02	3.6
Secondary Material	0.02	2.9E-06	1.1E03
Secondary Renewable Fuel	1.1E-02	6.7E-06	4.1E-04
Primary Renewable Material	3.8E-03	2.4E-03	3.1E-03
Primary Energy Renewable Not Feedstock	1.0	2.9E-04	6.5E-02
Primary Energy Renewable Total	1.0	2.7E-03	6.9E-02
Secondary Non-renewable Fuel	4.6E-03	7.4E-04	5.2E-04
Primary Energy Non-renewable Material	0.81	0.11	0.06
Primary Non-renewable Energy Not Feedstock	11	0.19	0.36
Primary Energy Non-renewable Total	10.6	0.30	0.42
Hazardous Waste Disposed	2.4E-04	3.7E-05	1.6E-05
Non-hazardous Waste Disposed	0.12	3.1E-04	5.5E-02
Radioactive Waste Disposed	3.7E-16	1.1E-31	1.5E-17
Components For Reuse	0	0	0
Material For Recycling	6.5E-03	6.5E-06	5.8E-03
Material For Energy Recovery	8.3E-05	2.3E-07	6.9E-06
Exported Energy Electrical	0	0	0
Exported Energy Thermal	0	0	0

Results Module B: Building Fabric and Operations

Table 2 shows B3 Repair results. Zero in B1 Use, B2 Maintain, B4 Replace, B5 Refurbish, B6 Energy Use, B7 Water Use

Table 2 B1 to B7 Impact & Inventory Results/Functional Unit

Result	B1	B2	B3	B4	B5	B6	B7
Climate Change biogenic	0	0	-4.3E-04	0	0	0	0
Climate Change luluc (landuse)	0	0	1.7E-07	0	0	0	0
Climate Change fossil	0	0	5.1E-02	0	0	0	0
Climate Change total	0	0	5.1E-02	0	0	0	0
Stratospheric Ozone Depletion	0	0	8.0E-10	0	0	0	0
Photochemical Ozone Creation	0	0	2.3E-04	0	0	0	0
Acidification Potential	0	0	1.0E-04	0	0	0	0
Eutrophication Freshwater	0	0	9.8E-09	0	0	0	0
Eutrophication Marine	0	0	2.4E-05	0	0	0	0
Eutrophication Terrestrial	0	0	5.8E-05	0	0	0	0
Fossil Depletion	0	0	2.4E-02	0	0	0	0
Mineral and Metal Depletion	0	0	1.7E-05	0	0	0	0
Water Scarcity Depletion	0	0	5.8E-04	0	0	0	0
Net Fresh Water Use	0	0	3.60	0	0	0	0
Secondary Material	0	0	1.1E-03	0	0	0	0
Secondary Renewable Fuel	0	0	4.1E-04	0	0	0	0
Primary Renewable Material	0	0	3.1E-03	0	0	0	0
Primary Energy Renewable Not Feedstock	0	0	6.5E-02	0	0	0	0
Primary Energy Renewable Total	0	0	6.9E-02	0	0	0	0
Secondary Non-renewable Fuel	0	0	5.2E-04	0	0	0	0
Primary Energy Non-renewable Material	0	0	5.7E-02	0	0	0	0
Primary Non-renewable Energy Not Feedstock	0	0	0.36	0	0	0	0
Primary Energy Non-renewable Total	0	0	0.42	0	0	0	0
Hazardous Waste Disposed	0	0	1.6E-05	0	0	0	0
Non-hazardous Waste Disposed	0	0	5.5E-02	0	0	0	0
Radioactive Waste Disposed	0	0	1.5E-17	0	0	0	0
Components For Reuse	0	0	0	0	0	0	0
Material For Recycling	0	0	5.8E-03	0	0	0	0
Material For Energy Recovery	0	0	6.9E-06	0	0	0	0
Exported Energy Electrical	0	0	0	0	0	0	0
Exported Energy Thermal	0	0	0	0	0	0	0

Results Module C: End-of-life

Table 3 shows results for C1 demolish, C2 Transport C4 Disposal. C3 Waste Processing has no flows.

Table 3 C1 to C4 Impact & Inventory Results/Functional Unit

Result	C1	C2	C3	C4
Climate Change biogenic	-1.0E-05	-1.0E-05	0	-7.8E-07
Climate Change luluc (landuse)	4.6E-11	1.4E-09	0	7.1E-10
Climate Change fossil	3.0E-06	6.0E-03	0	7.4E-03
Climate Change total	3E-06	6.0E-03	0	7.4E-03
Stratospheric Ozone Depletion	2.3E-13	1.1E-13	0	1.1E-13
Photochemical Ozone Creation	2.2E-08	6.0E-05	0	7.5E-05
Acidification Potential	1.4E-08	5.1E-06	0	2.0E-04
Eutrophication Freshwater	3.3E-13	3.1E-10	0	3.4E-10
Eutrophication Marine	4.2E-09	9.5E-07	0	1.2E-06
Eutrophication Terrestrial	7.4E-09	3.4E-06	0	3.8E-06
Fossil Depletion	2.1E-06	7.5E-03	0	9.0E-03
Mineral and Metal Depletion	3.8E-09	4.0E-06	0	4.9E-06
Water Scarcity Depletion	1.6E-07	1.4E-06	0	1.6E-06
Net Fresh Water Use	0.00	0.01	0	9.7E-03
Secondary Material	3.4E-07	2.2E-06	0	1.6E-06
Secondary Renewable Fuel	1.1E-07	5.1E-06	0	4.7E-06
Primary Renewable Material	1.4E-07	1.6E-03	0	2.0E-04
Primary Energy Renewable Not Feedstock	1.5E-05	2.0E-04	0	2.0E-04
Primary Energy Renewable Total	1.5E-05	1.8E-03	0	1.9E-03
Secondary Non-renewable Fuel	1.4E-08	4.8E-04	0	5.1E-04
Primary Energy Non-renewable Material	2.4E-06	0.04	0	0.04
Primary Non-renewable Energy Not Feedstock	4.3E-05	0.06	0	0.08
Primary Energy Non-renewable Total	4.6E-05	0.10	0	0.12
Hazardous Waste Disposed	7.1E-10	1.2E-05	0	1.5E-05
Non-hazardous Waste Disposed	1.4E-06	9.7E-05	0	1.0
Radioactive Waste Disposed	4.4E-21	8.5E-32	0	7.5E-32
Components For Reuse	0	0	0	0
Material For Recycling	1.5E-08	4.6E-06	0	4.0E-06
Material For Energy Recovery	2.9E-10	1.5E-07	0	1.6E-07
Exported Energy Electrical	0	0	0	0
Exported Energy Thermal	0	0	0	0

Results Module D: Beyond System Boundaries

Table 3 has results for benefit and loads in D1 reuse and D2 recovery. D3 recycling has no flows.

Table 3 D1 to D3 Impact & Inventory Results/Functional Unit

Result	D1	D2	D3
Climate Change biogenic	-2.0E-04	-1.9E-4	0
Climate Change luluc (landuse)	1.7E-07	2.4E-09	0
Climate Change fossil	4.8E-02	0	0
Climate Change total	4.8E-02	0	0
Stratospheric Ozone Depletion	8.2E-10	5.9E-13	0
Photochemical Ozone Creation	2.3E-04	1.2E-06	0
Acidification Potential	1.0E-04	5.3E-07	0
Eutrophication Freshwater	2.2E-09	1.2E-10	0
Eutrophication Marine	2.4E-05	9.4E-08	0
Eutrophication Terrestrial	5.8E-05	6.9E-07	0
Fossil Depletion	2.4E-02	1.7E-04	0
Mineral and Metal Depletion	1.8E-05	5.8E-08	0
Water Scarcity Depletion	6.0E-04	1.8E-05	0
Net Fresh Water Use	3.7	0.11	0
Secondary Material	1.1E-03	0	0
Secondary Renewable Fuel	3.8E-04	4.3E-05	0
Primary Renewable Material	4.4E-05	3.0E-05	0
Primary Energy Renewable Not Feedstock	6.0E-02	1.4E-04	0
Primary Energy Renewable Total	6.0E-02	1.7E-04	0
Secondary Non-renewable Fuel	2.9E-04	7.7E-06	0
Primary Energy Non-renewable Material	4.5E-02	0	0
Primary Non-renewable Energy Not Feedstock	0.35	3.1E-03	0
Primary Energy Non-renewable Total	0.42	3.1E-03	0
Hazardous Waste Disposed	1.4E-05	1.9E-07	0
Non-hazardous Waste Disposed	7.3E-03	2.0E-05	0
Radioactive Waste Disposed	1.6E-17	4.9E-21	0
Components For Reuse	0	0	0
Material For Recycling	1.9E-04	1.6E-05	0
Material For Energy Recovery	7.3E-06	6.5E-09	0
Exported Energy Electrical	0	0	0
Exported Energy Thermal	0	0	0

Interpretation Cradle to Gate A1 to A3.

The first interpretation section discusses product results cradle to gate A1 to A3.

Figure 3 charts such mass versus EE/kg. It shows highest sensitivity Calcium aluminate cement and least sensitivity to sand content.

The Calcium aluminate cement and proprietary base mix were significantly more energy intensive than others.

Figure 4 charts mass versus GWP/kg product. It shows highest sensitivity to calcium aluminate cement and least sensitivity to sand content.

Calcium aluminate, base mix, hydrated lime and Portland cement components were significantly more CO_{2e} intensive than the sand.





Figure 3 Mass Share Vs Embodied Energy MJ//kg A1-3 **Figure 4 Mass Share Vs CO_{2e} kg/kg A1-3**

Interpretation Cradle to Grave and Beyond the System Boundary A1 to D3.

The second interpretation section discusses product results cradle to grave and beyond A1 to D3. With product lasting beyond 60-years.

Figure 5 shows highest GWP A1-A3 and insignificant A4 to C4.



Figure 5 GWP A1 to D3/kg Functional Unit

References

[1] EN 15804:2012+A2:2019 Sustainability of construction works - Environmental product declarations - Core rules for the product category of construction products.

[2] GreenTag™ 2023 EPD Program, Product Category Rules https://www.globalgreentag.com/epd-program.html Sub-PCR UCM:2023 Unreinforced Concrete Mixtures and Additives

[3] ISO 14025:2010 Environmental labels and declarations – Type III – environmental declarations - Principles and procedures.

[4] ISO14044:2006 Environmental management – Life cycle assessment – Requirements and guidelines.

[5] ISO 15686-2:2012 Buildings and constructed assets - Service life planning - Part 2: Service life prediction procedures.

[6] ISO 15686-8:2008 Buildings and constructed assets - Service-life planning - Part 8: Reference service life and service-life estimation.

[7] IPCC 2013, Global Warming Potential 100-year, IPCC Fifth Assessment Report Climate Change.

[8] WMO 2014, Ozone Depletion Potentials for Steady-state, Scientific Assessment of Ozone Depletion: 2014, Global Ozone Research and Monitoring Project Report No. 55, 2014.

[9] Van Zelm, R., Huijbregts, M., Hollander, H., Jaarsveld, H., Sauter, F., Struijs, J., Wijnen, H., Van de meent, D. 2008, European characterization factors for human health damage of PM10 and ozone in life cycle impact assessment, J O Atmospheric Environment 42(3):441-453, as applied in ReCiPe LOTOS-EUROS. DOI: 10.1016/j.atmosenv.2007.09.072

[10] Seppälä, J., Posch, M., Johansson, M. and Hettelingh, J-P. 2006 Country-dependent Characterisation Factors for Acidification and Terrestrial Eutrophication Based on Accumulated Exceedance as an Impact Category Indicator, T Int J O LCA 11(6):403-416 Nov 2006 DOI:10.1065/lca2005.06.215

[11] Posch, M., Seppälä, J., Hettelingh, J-P., and Johansson, M., (2008) The role of atmospheric dispersion models and ecosystem sensitivity in the determination of characterisation factors for acidifying and eutrophying emissions in LCIA, Sept 2008, I J of Life Cycle Assessment 13(6):477-486., DOI:10.1007/s11367-008-0025-9

[12] Struijs, J., Beusen, A., van Jaarsveld, H. & Huijbregts, M.A.J. (2009b). Aquatic Eutrophication. Ch 6 in: Goedkoop, M., Heijungs, R., Huijbregts, M.A.J., De Schryver, A., Struijs, J., Van Zelm, R. (2009). ReCiPe 2008 A life cycle impact assessment method which comprises harmonised category indicators at the midpoint and the endpoint level. Report I: Characterisation factors, 1st Ed.

[13] CML–IA V4.1 LCA methodology, 2002, October 2012, CML University of Leiden, Netherlands.

[14] Guinée et al., 2002, and van Oers et al., 2002 CML LCA methodology 2002a, Institute of Environmental Sciences (CML), Faculty of Science, University of Leiden, Netherlands.

[15] Boulay, A-M., Bare, J., Benini, L., Berger, M., Lathuilliere, M., Manzardo, A., Margni, M., Motoshita, M., Núñez, M., Pastor, A., Ridoutt, B., Oki, T., Worbe, S., Pfister, S. (2018). The WULCA consensus characterization model for water scarcity footprints: assessing impacts of water consumption based on available water remaining (AWARE). I J of LCA. 23. 1-11. 10.1007/s11367-017-1333-8.

Bibliography

Ciroth A., Hildenbrand J., Zamagni A. & Foster C., 2015, Data Review Criteria. Annex A: LCI Dataset Review Criteria, 10.13140/RG.2.1.2383.4485 UN EP Life Cycle Initiative

EN ISO 14024:2000, Environmental labels and declarations - Type I environmental labelling -Principles and procedures (ISO 14024:1999).

EN ISO 14040:2006, Environmental management - Life cycle assessment - Principles and framework (ISO14040:2006).

EN 15643-1:2010, Sustainability of construction works - Sustainability assessment of buildings - Part 1: General framework.

EN 15643-2, Sustainability of construction works - Assessment of buildings - Part 2: Framework for the assessment of environmental performance.

EN 16449, Wood and wood-based products - Calculation of the biogenic carbon content of wood and conversion to carbon dioxide.

ISO 21930:2007 Sustainability in building construction - Environmental declaration of building products.

ISO 21931-1:2010, Sustainability in building construction - Framework for methods of assessment of the environmental performance of construction works - Part 1: Buildings.